



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

defining the range of animals whose range one knows, in terms of the ranges of plants and animals whose range one does not know.

Lutz' method of "Geographic Average" (1921, *Amer. Mus. Novitates*, 5) involves considerable calculation, which brings to light with great effort the astounding information that the "mid-range" of a northern species is north of the "mid-range" of a southern species.

He voices considerable discontent with the zones as being indefinite and indistinct. Whereupon he proceeds to define anew the undefinable and to redistinguish the indistinguishable.

Inasmuch as all sharp changes in fauna are dependent on similarly marked changes in topography, it would seem reasonable to adopt physiographic regions, instead of zones, "major environments," and "geographic averages."

For the United States this has been made remarkably easy by the publication of "Physiographic Divisions of the United States," by Nevin S. Fenneman, *Ann. Ass. Amer. Geog.*, VI, 1917, which is the result of long continued work on the part of a committee of the association.

The lines of these divisions in many cases agree with the zones and with the plant formations. This is to be expected for the ranges of many animals should logically stop at a physiographic break, or at a line of sudden change in temperature or moisture, and the former usually carries with it the two latter.

Furthermore the use of these physiographic regions clears the way for intelligent association of animal and plant ranges with the geology and soil conditions, and brings to light routes of dispersal.

For instance, this system makes clear the presence of the Connecticut valley and the Hudson-Champlain trough, in which many southern forms extend north, but to draw these on the map as Upper Austral, indicating faunal identity with Piedmont Virginia (also Upper Austral) is sheer falsehood.

The beauty of this method, however, is its ease of application. These regions can be *seen*, their boundaries can be *seen*, and there is seldom any doubt as to which region one is in or from which specimens came.

Of course, all animals do not respect physiographic boundaries, any more than all animals respect the lines drawn on maps to represent faunal zones.

None of our methods of indicating general distribution is perfect. None ever can be. Zoogeographers might well give some consideration to the results of the "new geography," and afford themselves some justification for their title.

EMMETT REID DUNN

SMITH COLLEGE

CLINKERTILL, A NEW METAMORPHIC ROCK

IN view of the interest attached to the study and identification of tillite a note about a rather unusual form of till may be of interest. This occurrence has never been described so far as the writer is aware.

Capping the northern bluffs bordering the Missouri river in portions of sections 16, 17 and 18, Township 154 west, Range 95 north, Williams County, North Dakota, is a typical boulder-clay. This till lies on the truncated edges of the gently easterly dipping, lignite bearing, Fort Union Beds. In the sections mentioned above a heavy bed of lignite, here known as the Williston bed¹ immediately underlies the drift. Here as elsewhere the Williston bed has burned back from the outerop baking the overlying till in places to a thickness of 30 feet or 40 feet. The clinkertill so formed varies in color from salmon-pink to dark brown, brick-red being the common color. In selected spots the material is fused to a porous, scoria-like mass but in most part is only slightly indurated by the heat. This baked till so formed resists weathering and removal and forms the capping of steep bluffs and buttes or mesas.

The most common pebbles found in this clinkertill are granite and limestone. The granite pebbles appear to be unaltered by the heat and the limestone pebbles but slightly as they effervesce freely in hydrochloric acid.

L. P. DOVE

NORTH DAKOTA GEOLOGICAL SURVEY

¹ Collier, A. J., "The Nesson Anticline," Bulletin 691 G., U. S. G. S., p. 213.